

## Applied evolutionary ecology of insects of the subfamily Bruchinae (Coleoptera: Chrysomelidae)

Midori TUDA\*

Institute of Biological Control, Faculty of Agriculture, Kyushu University, Fukuoka 812–8581, Japan

(Received 17 October 2006; Accepted 19 January 2007)

### Abstract

Bean beetles of the subfamily Bruchinae (formerly, the family Bruchidae) include notorious pests of stored legumes, *Callosobruchus*, *Caryedon*, *Acanthoscelides* and *Zabrotes* that are able to feed and reproduce on dried beans and peas. Here, I review recent findings on the ecology, phylogeny, invasion and evolution in the bean beetles, based on field investigation of host plants and molecular studies. Possible future application of the new knowledge to weed and pest control is proposed, such as potential utility of the seed predators for modest control of beneficial yet invasive ('conflict') plants and new control methodology of pest bean beetles.

**Key words:** Bruchidae; seed predators; stored product pests; biological weed control; molecular phylogeny

### INTRODUCTION

Beetles of the subfamily Bruchinae (Coleoptera: Chrysomelidae) are specialized internal feeders of bean family seeds (Fabaceae). They are frequently and erroneously referred to as the bean 'weevils' in spite of their affinity to the leaf beetles (Chrysomelidae). Here, I use the common name, the bean 'beetle' (or the seed 'beetle') to avoid taxonomic confusion. I also follow the recent taxonomic consideration that the bean beetle is not unique enough as a family but as a subfamily of Chrysomelidae (Lingafelter and Pakaluk, 1997).

The Bruchinae consist of about 1,700 species (Johnson et al., 2004). Although not very high in proportion of species, some of the bean beetles are notorious pests of stored beans with the highest intrinsic rate of increase among stored-products pests (Imura, 1990). They include several species of *Callosobruchus*, *Acanthoscelides* and *Zabrotes* that infest beans (Tribe Phaseoleae: *Vigna*, *Phaseolus*, *Glycine*, etc.) and *Caryedon* that feeds on peanuts or groundnuts (*Arachis hypogaea*) (Southgate, 1979; Johnson, 1981). *Bruchus* species that infest broad beans and peas (Fabeae: *Vicia* and *Pisum*) of economic importance may also be categorized as

stored-bean pests. Strictly speaking, however, the genus *Bruchus* is different from other stored-bean pests in that it does not reproduce on dried hardened beans/peas and thus cannot cause further damage in storage.

### PHYLOGENY

Recent molecular studies support that the Bruchinae are closely related to the frog-legged beetle subfamily, Sagrinae, within the family Chrysomelidae (Farrell and Scqueira, 2004). Within the Bruchinae, the tribes Amblycerini and Pachymerini are thought to be paraphyletic to a major tribe Bruchini (Farrell, 1998; Table 1). Although recent evidence supports the previously proposed phylogeny at the higher taxonomic levels, an intriguing new view of bean beetle phylogeny at the lower levels has been proposed. Kergoat et al. (2005a, b) supported the previously suggested view that *Acanthoscelides* and *Bruchidius* are polyphyletic groups (Johnson, 1981; Borowicz, 1987). However, some monophyletic groups of *Acanthoscelides* may be more closely related to those of *Bruchidius* than the other congeneric groups (Alvarez et al., 2006; Tuda, Kergoat, Kato, Ito, Bu-

Table 1. Taxonomic position and composition of Bruchinae

Family	Subfamily	Tribe
Chrysomelidae		
	Bruchinae	
	Amblicerini	3 genera ( <i>Spermophagus</i> , <b><i>Zabrotes</i></b> )
	Bruchini	46 genera ( <i>Acanthoscelides</i> , <i>Bruchidius</i> , <b><i>Bruchus</i></b> , <i>Callosobruchus</i> , <i>Mimosestes</i> )
	Eubaptini	—1 genus ( <i>Eubaptus</i> )
	Kytorhinini	—1 genus ( <i>Kytorhinus</i> )
	Pachymerini	—12 genera ( <b><i>Caryedon</i></b> )
	Rhacbini	—1 genus ( <i>Rhacbus</i> )

The genera in bold include pest species.

ranapanichpan, Szentesi and Jermy, unpublished). These molecular studies also indicate potential problems about relative phylogenetic positions among other recently described genera. For example, a recent molecular study found *Tuberculo-bruchus* to be polyphyletic (Kergoat, 2005b), indicating a plesiomorphic trait is used for the generic key and revision of the genus using genital morphology may be necessary. Data on recently described species, synonymies and host plants have been actively accumulated from different biogeographic regions of Asia and Africa that would contribute to the understanding of evolutionary diversity of the Old World (e.g., Arora, 1977, 1980; Egorov and Ter-Minassian, 1983; Morimoto, 1990; Anton et al., 1997; Anton, 1999, 2000; Kingsolver, 1999; Tuda, 2003; Johnson et al., 2004; Tuda and Morimoto, 2004; Tuda et al., 2005) and the New World (e.g., Johnson, 1970; Romero and Johnson, 2000) bean beetles (see also Zacher, 1952; Udayagiri and Wadhi, 1989 for overviews).

## ECOLOGY

In most bean beetles, female adults deposit eggs on pods and/or seeds, hatched larvae burrow into seeds and emerge out as adults (Southgate, 1979). Adults lay eggs singly and larvae feed only single seeds (Center and Johnson, 1974). Exceptions to the typical life history are *Conicobruchus* (Prevett, 1967), *Merobruchus* (Johnson, 1967) and *Sennius* (Center and Johnson, 1972), which feed on several seeds in a single pod and *Bruchidius*, which feeds on multiple pods in single *Trifolium* inflorescences

(Balachowsky, 1962; Tuda, pers. obs.). In both cases, the host seeds are too small to support bean beetle development into full adult. Finally, as an exception to the general pattern of pupating within seeds, the larvae of several species of *Caryedon* emerge out of seeds to form cocoons either on pods or on the ground for pupation (Center and Johnson, 1974; Southgate, 1979; Tuda, pers. obs.).

Batch egg laying is known only for a limited number of species such as *Pseudopachymerina spinipes* (Teran, 1962), *Caryedon fasciatus* (Prevett, 1966), *Caryoborus serripes* (Delobel et al., 1995a) and *Stator beali* (Nilsson and Johnson, 1993; Fox and Mousscau, 1995). The host pods and seeds of these beetles are hard and relatively large (Janzen, 1971; Moreno-Casasola et al., 1994; Delobel et al., 1995a; Delgado et al., 1997). In *Callosobruchus*, such egg clustering had previously not been observed but it was recently found in a non-pest species, which develops in a large, hard seeded legume (Tuda and Buranapanichpan, unpublished). These examples imply that egg laying pattern is not phylogenetically constrained but may respond to selection imposed by host plant characteristics (the size and/or hardness of pods and seeds), parasitoid pressure and abiotic factors (Janzen, 1971; Mitchell, 1977; Delobel et al., 1995a).

Typical hosts of bean beetles are legumes (Fabaceae) (84% of the known hosts; Johnson, 1970) but the palm (Arecaceae), morning glory (Convolvulaceae), mallow (Malvaceae) and about 30 other families are also used as hosts (Southgate, 1979; Johnson, 1981). Most bean beetles are either monophagous or oligophagous; their host range is limited to restricted plant taxa, usually subtribes and tribes at most (Jermy and Szentesi, 2003; Delobel and Delobel, 2003; Tuda et al., 2005; Kergoat et al., 2007a, b), although there are a few generalists that utilize different subfamilies of legumes.

Evolutionary relation between bean beetles and their host plants has been actively discussed as a model of plant-insect evolution. Proximate mechanisms for host-plant shifts are probably associated with chemical (Janzen, 1969), morphological (Janzen, 1969; Szentesi and Jermy, 1995) and geographical (Johnson and Siemens, 1991) proximity between the current and potential hosts. Delobel and Delobel (2006) suggest that the speciation in European bean beetles is sequential evolution

(sensu Jermy, 1984) or sequential radiation (sensu Abrahamson et al., 2003) that follows plant diversification, which contrasts to the reciprocal coevolution model that assumes plant evolution in response to insect evolution against plant defensive traits (sensu Ehrlich and Raven, 1964; Futuyma, 1983; Becerra, 2003).

## EVOLUTION OF STORED PRODUCT PESTS

Evolution of stored bean pests is likely a two-step process. The first is preadaptation to utilize dried hard seeds (Watanabe, 1985) and the second is the evolution of dispersal, reproductive and competitive polymorphism to adapt to cultivated seeds and storage environments (Utida, 1954, 1981; Caswell, 1960; Nakamura, 1966; Sano, 1967; Ouedraogo and Huignard, 1981; Messina, 1984; Toquenaga, 1990; Tuda, 1997, 1998; Tuda and Iwasa, 1998; Takano et al., 2001). Hence, an ability to feed on dried mature host seeds is an inevitable prerequisite for bruchids to become pests. This requires first, female adults to deposit eggs on/near dried mature seeds and second, hatched larvae to burrow into the hard seeds to feed on.

What is it then that promotes evolution of utilization of dry, hard seeds? Using a comparative approach to study *Callosobruchus* pest and non-pest species, the potential effects of climate, host plants, phylogeny and endosymbiotic bacterium were examined (Tuda et al., 2006). Long dry season was shown to be the most important factor for evolution towards preadaptive stage to become stored bean pests (i.e., ability to use dry hard beans) when evolutionary history was accounted for (Tuda et al., 2006). Phylogenetic history of *Callosobruchus* and idiosyncrasy of their host plants also had significant effects on the evolution towards stored bean pests (Tuda et al., 2006). In fu-

ture studies, hardness of dried seeds will be one of the foci of research because the hardness per se can serve as a deterrent against seed predators, including bean beetles (Janzen, 1977; Southgate, 1979; Kitch et al., 1991; Dongre et al., 1993). The loss of toxic chemicals during post-maturity drying processes may also increase survival of such granivores, which promotes host range expansion.

## INVASION

With human aid, dispersal and invasion of living organisms in recent centuries are occurring at a speed that has been unattainable with plate tectonics and glacial dynamics. In Japan, examples of bean beetle invaders are *Bruchus rufimanus* and *Bruchus pisorum*, the pests of broad beans and peas, respectively (Table 2). *Bruchus lori* has also been considered to be an alien species but our recent finding indicates the current status as an introduced species may require reconsideration, as described later in this section. We found two additional bean beetle species have invaded Japan (Tuda et al., 2001; Tuda, Tateishi, Niyomdham, Morimoto, Chen, Zhu, Zhang, Murugan, Chou and Johnson, unpublished).

One recent invader is a bean beetle *Acanthoscelides pallidipennis* (= *A. collusus*) that feeds on a fast-growing, nitrogen-fixing legume, Indigobush (or False indigo; Itachi-hagi), *Amorpha fruticosa*. The legume had invaded from North America and established in England, Europe and East Asia except Japan by the mid 20th century (Szentesi, 1999; Tuda et al., 2001). In Japan, the bean beetle was first found at two locations in the central and southwestern parts of the country in 1997 (Tuda et al., 2001) and later in several areas in the southwestern Japan as well (Ishihara, unpublished). A molecular analysis of *A. pallidipennis*

Table 2. Bean beetle invasion to Japan

Bean beetle	Time of invasion	Source population
<i>Bruchus pisorum</i> <sup>a</sup>	1888	USA
<i>Bruchus rufimanus</i> <sup>a</sup>	1921	England/Europe
<i>Callosobruchus maculatus</i> <sup>b</sup>	1950	?
<i>Acanthoscelides pallidipennis</i> <sup>c</sup>	1970s-1997	Korea and China
<i>Acanthoscelides macrophthalmus</i> <sup>d</sup>	recent (2000)	?

<sup>a</sup>Yoshida, 1990; <sup>b</sup>Morimoto and Kiritani, 1995; <sup>c</sup>Tuda et al., 2001; <sup>d</sup>Tuda, Tateishi, Niyomdham, Morimoto, Chen, Zhu, Zhang, Murugan, Chou and Johnson, unpublished.

from four States of the USA and from East Asia revealed that Japanese populations consist of several unrelated haplotypes that are shared by Chinese and Korean populations and individuals intercepted by import inspection at Japanese seaport quarantines (Tuda, Ishihara, Wasano, Morimoto, Paik, Houck and Johnson, unpublished). This molecular evidence supports our hypothesis (Tuda et al., 2001) that *A. pallidipennis* established in Japan was transported from the introduced populations in nearby East Asian countries. Indeed, from these regions, the host seeds (*A. fruticosa*) have been imported for erosion control since the 1970s (Tuda et al., 2001).

The other invader is *Acanthoscelides macrophthalmus* (Tuda, Tateishi, Niyomdham, Morimoto, Chen, Zhu, Zhang, Murugan, Chou and Johnson, unpublished). It is a Neotropical species that feeds on Wild tamarind, or Gin-nemu, *Leucaena leucocephala*. *Leucaena leucocephala* is a fast-growing nitrogen-fixing leguminous tree that is cultivated for fodder (Elharith et al., 1980), green manure (Chagas, 1981), reforestation, windbreak, fuel, pulp and erosion control (Kondo et al., 1987; Satake et al., 1989). Because *A. macrophthalmus* is specialized to *Leucaena* species (e.g., Johnson, 1979; Hughes and Johnson, 1996; Delobel and Johnson, 1998), *L. leucocephala* is the only *Leucaena* species that has been introduced to Japan, and airborne long-distance dispersal is highly unlikely for bean beetles (Tuda et al., 2001), there is little doubt that *A. macrophthalmus* has been introduced with the host seeds.

Accelerated long-range human transportation promotes unexpected dispersal of animals and plants that eventually become pests in the areas where they are introduced. As a consequence, it is sometimes difficult to determine geographic origins of widely distributed organisms. Whether local populations are native or of relatively recent introduction from foreign populations can be estimated by statistics based on population genetics theory if DNA sequence or restriction site data are available (e.g., Templeton et al., 1992; Templeton, 1998). Our recent findings regarding Japanese populations of *Bruchus loti* may be a good example. Palearctic fauna and flora share many species including *B. loti*. The Japanese populations of *B. loti* are currently listed as an alien species that would become a target of eradication. We found that hap-

lotypes found in the Japanese populations are clearly different from those in European populations, which indicates that Japanese *B. loti* is not introduced from Europe and likely originated from the Far East including Japan (Tuda, Ichita and Kergoat, unpublished). The genetic uniqueness of the Japanese population indicates that *B. loti* should perhaps be conserved in this country. The status of Japanese *Callosobruchus chinensis* is also controversial. Our molecular study shows that populations of *C. chinensis* in agricultural habitats are genetically homogeneous, whereas those from natural habitats are heterogeneous, indicating a bottleneck event caused by human agricultural practice and transportation (Tuda et al., 2004).

## APPLICATION TO BIOLOGICAL CONTROL

**As natural enemy of conflict plants.** A control agent should ideally be a specialist feeding only on targets of control and not harm beneficial organisms (Sweetman, 1936; Huffaker, 1964). Internal feeders ('endophages') tend to have narrower diet breadth than external feeders ('exophages') (Lewinsohn, 1991; Frenzel and Brandl, 1998) and are probably more suitable as biological control agents. Indeed, bean beetles have been used as biological control agents of weedy plants and proven to be effective, to some extent, once established in the release areas (Table 3; c.f., Julien, 1992).

I propose two bean beetle species that have been accidentally introduced and can be used for biological control of new leguminous weeds in Japan. The first is *A. pallidipennis*. The narrow diet range of *A. pallidipennis* and the absence of congeneric relatives of Indigobush and naturally associated parasitoids of the beetle are preferable features as a control agent, in contrast with the case of *Bruchidius villosus* that attacked not only the target but also non-target (but non-indigenous) congeners after its introduction to New Zealand (Paynter et al., 2004). In this respect, *A. pallidipennis* is suitable as an agent that controls escape of *Amorpha fruticosa* from cultivation by seed dispersal. The suitability of *A. pallidipennis* as a control agent has also been pointed out in the native distribution area of Indigobush (Rogers and Garrison, 1975). The bean beetle would be even more suitable if it would be applied for control in introduced areas where its naturally associated enemies (i.e., parasitoids,

Table 3. Bean beetles used as biological control agents of weeds (modified from Julien, 1992; ARC-PPRI, 2003)

Control agent	Target weed	Location of release
<i>Acanthoscelides macrophthalmus</i>	<i>Leucaena leucocephala</i>	South Africa
<i>Acanthoscelides puniceus</i>	<i>Mimosa pigra</i>	Australia, Thailand, Vietnam (Malaysia, Myanmar)
<i>Acanthoscelides quadridentatus</i>	<i>Mimosa pigra</i>	Australia, Thailand, Vietnam (Malaysia, Myanmar, USA)
<i>Algarobius bottimeri</i>	<i>Prosopis glandulosa</i>	South Africa
<i>Algarobius prosopis</i>	<i>Prosopis velutina</i>	South Africa
<i>Bruchidius villosus</i>	<i>Cytisus scoparius</i>	New Zealand
<i>Bruchidius sahlbergi</i>	<i>Acacia nilotica</i>	Australia
<i>Sulcobruchus subsuturalis</i>	<i>Caesalpinia decapetala</i>	South Africa

Locations in parentheses are due to natural spread of released agents.

predators, parasites, and pathogens) and competitors (e.g., *Acanthoscelides submuticus*) are rare or absent (Tuda et al., 2001). *Acanthoscelides pallidipennis* feeds only on *Amorpha* and related genera (Amorphaeae: *Errazurizia* and *Parryella*) that distribute naturally only in North America (Center and Johnson, 1974).

The second is *A. macrophthalmus*. *Leucaena leucocephala*, initially introduced as a beneficial tree in the 19th century, escaped from cultivation by seed dispersal and has become weedy in tropical regions of Japan and other introduced areas (e.g., Smith, 1985; Henderson, 2001; Wu et al., 2003). Its seed predator, *A. macrophthalmus*, satisfies the above criteria as a control agent in Asia, i.e., narrow host range and scarcity of parasitoids. In fact, this bean beetle has already been deliberately introduced to South Africa for the control of *L. leucocephala* (ARC-PPRI, 2003; Olckers, 2004; Table 3). Nevertheless, there remains a risk that the control agent could easily accumulate local generalist parasitoids and in the long term the control effect could decrease as observed in West Africa (Delobel and Johnson, 1998).

**Control of bean beetles.** Fumigation has been widely applied to stored products as an effective method of control of stored products pests. However, methyl bromide, used for fumigation of grain legumes (Yoneda et al., 1990) has been recognized as one of the chemicals that deplete the stratospheric ozone layer and, hence, was called for its controlled use by the Montreal Protocol (UNEP, 2000).

Alternatively, parasitoids (e.g., a trichogrammatid egg parasitoid *Uscana*, larval-pupal parasitoids, pteromalid *Dinarmus*, *Anisopteromalus*, an eupelmid *Eupelmus*, and a braconid *Heterospilus*;

e.g., Southgate, 1979; Steffan, 1981; Fujii and Wai, 1990; Mitsunaga and Fujii, 1999; Schmale et al., 2001; Tuda et al., 2001; Kobayashi et al., 2003; Jaloux et al., 2004; Tuda and Shimada, 2005; Wu et al., 2005; Vamosi, Hollander and Tuda, unpublished) and plant extracts (e.g., Lambert, 1985; Rahman, 1990) have been actively explored for biological control (see Fujii et al., 1990; Huis, 1991 for review). In addition to biological and botanical control, controlled temperature (e.g., Rahman, 1990), mechanical control (e.g., Quentin et al., 1991), controlled atmosphere (e.g., Oosthuizen and Schmidh, 1942) and radiation (e.g., Kiyoku and Tsukuda, 1968; Hossain et al., 1972; Reddy et al., 2006) have been proposed as effective control methodology (see also Highley et al., 1994). I suggest an endosymbiotic bacterium, *Wolbachia*, from bean beetle pests and its functional relation with the host (Kondo et al., 2002) may be applied to biological control of stored product pests. When the above-mentioned multiple methods are used in combination, the possibility of one agent negating the other must be taken into account (e.g., Boeke et al., 2003).

**Identification of pest species.** External and internal morphology of the adult stage is used for identification to species. Most recently male genital traits for *Callosobruchus* were compared and summarized by Tuda et al. (2006). Compared to adult morphology, egg and larval morphology is less well studied because they are not readily available (but see Pfaffenberger and Johnson, 1976; Arora, 1978; Delobel et al., 1995b), which makes identification based on pre-adult stages difficult. In contrast to the differential accessibility among developmental stages of morphological characters, molecular characters can be more stable and useful as

keys for identification. Using the PCR-RFLP technique, which is relatively easy and inexpensive, species of all developmental stages of important stored bean pests in the genus *Callosobruchus* have been successfully distinguished (Tuda et al., 1995).

## CONCLUSION

Bean beetles, as other phytophagous insects, exhibit significant conservatism in host utilization. Feeding on dried seeds not only enables repeated generations but also broadens the diet of some bean beetle pests. Our study based on molecular phylogenetics indicates this preadaptation to use dry, hard beans that precedes human store of beans has been selected for by arid habitat climate (Tuda et al., 2006). Recent investigations of Asian bean beetle fauna have revealed diversity of the species and ecology of the seed predators. This also led to the finding of invasions of alien bean beetles (Tuda et al., 2001; Tuda et al., unpublished). Invasion routes estimated by our molecular approach indicated the invasion of the bean beetle was associated with imported alien host seeds. I propose the recent bean beetle invaders may be used for modest control of the seed dispersal of beneficial but invasive alien plants ('conflict plants' (Neser, 1994)).

## ACKNOWLEDGEMENTS

I am grateful to the Japanese Society of Applied Entomology and Zoology for the honorable award and the opportunity to write this review. Thanks are also due to Alex Delobel and Steven Vamost for their valuable comments on the manuscript. This study was supported partly by the Fujiwara Natural History Foundation, the Sumitomo Foundation, and Grant-in-Aids for International Scientific Research (Field Research 09041145) and for Scientific Research (A) (08304049, 15208007), (B) (14405003, 17405005) and for Young Scientists (B) (15770011) from MEXT. This paper is dedicated to my collaborators, the late Prof. Shwu-Bin Horng for his work on behavioral ecology of *Callosobruchus* and the late Dr. Liang-Yih Chou for his contribution to applied entomology.

## REFERENCES

Abrahamson, W. G., C. P. Blair, M. D. Eubanks and S. A. Morehead (2003) Sequential radiation of unrelated organisms: the gall fly *Eurosta solidaginis* and the tumbling flower beetle *Mordellistena convicta*. *J. Evol. Biol.* 16: 781–789.

Alvarez, N., J. R. Napoles, K.-W. Anton, B. Benrey and M. Hossaert-McKey (2006) Phylogenetic relationships in the Neotropical bruchid genus *Acanthoscelides* (Bruchinae, Bruchidae, Coleoptera). *J. Zool. Syst. Evol. Res.*

44: 63–74.

Anton, K.-W. (1999) Revision of the genus *Sulcobruchus* Chujo 1937 and description of *Parasulcobruchus* nov. gen. (Coleoptera, Bruchidae, Bruchinae). *Lin. Biol. Beitr.* 31: 629–650.

Anton, K.-W. (2000) Five new species of the *Callosobruchus chinensis* group from the Oriental Region and Australia (Coleoptera: Bruchidae: Bruchinae). *Genus* 11: 13–28.

Anton, K.-W., J. Halperin and M. Calderon (1997) An annotated list of the Bruchidae (Coleoptera) of Israel and adjacent areas. *Israel J. Entomol.* 31: 59–96.

ARC-PPRI (2003) Releases of Biological Control Agents against Weeds in South Africa. <http://155.240.199.39.institutes/ppri/main/divisions/weedsdiv/releases.htm>.

Arora, G. L. (1977) Taxonomy of Bruchidae (Coleoptera) of Northwest India. Part I. Adults. *Orient. Insect Suppl.* 7: 1–132.

Arora, G. L. (1978) Taxonomy of Bruchidae (Coleoptera) of Northwest India. Part II. Larvae. *Orient. Insect Suppl.* 8: 1–48.

Arora, G. L. (1980) *A Study of the Biology and Taxonomy of the Genus Bruchidius (Coleoptera: Bruchidae) from India. Final Tech. Rept (1974–1979) U.S. PL-480 Res. Proj. A7-ENF-103.* Department of Zoology, Punjab University, Chandigarh. 96 pp.

Balachowsky, A. S. (1962) *Entomologie Appliquée à l'Agriculture Tome I, Coléoptères*. Vol. 1. Masson et Cie Éditeurs, Paris. 564 pp.

Becerra, J. X. (2003) Synchronous coadaptation in an ancient case of herbivory. *Proc. Natl. Acad. Sci. USA* 100: 12804–12807.

Boeke, S. J., A. A. C. Sinzogan, R. P. de Almeida, P. W. M. de Boer, G. Jeong, D. K. Kossou and J. J. A. van Loon (2003) Side-effects of cowpea treatment with botanical insecticides on two parasitoids of *Callosobruchus maculatus*. *Entomol. Exp. Appl.* 108: 43–51.

Borowiec, L. (1987) The genera of seed-beetles (Coleoptera, Bruchidae). *Polsk. Pismo Entomol.* 57: 3–207.

Caswell, G. H. (1960) Observations on an abnormal form of *Callosobruchus maculatus* (F.). *Bull. Entomol. Res.* 50: 671–680.

Center, T. D. and C. D. Johnson (1972) Comparative life histories of *Sennius*. *Environ. Entomol.* 2: 669–672.

Center, T. D. and C. D. Johnson (1974) Coevolution of some seed beetles (Coleoptera: Bruchidae) and their hosts. *Ecology* 55: 1096–1103.

Chagas, J. M. (1981) *Leucaena leucocephala* as a green manure for bean growing in cerrado soil. *Pesq. Agr. Bras.* 16: 809–814.

Delgado, C., G. Couturier and A. Delobel (1997) Oviposition of seed-beetle *Caryoborus serripes* (Sturm) (Coleoptera: Bruchidae) on palm (*Astrocaryum chambira*) fruits under natural conditions in Peru. *Ann. Soc. Entomol. Fran.* 33: 405–409.

Delobel, A. and B. Delobel (2003) Les plantes hôtes des bruches (Coleoptera: Bruchidae) de la faune de France, une analyse critique. *Bull. Mens. Soc. Linn. Lyon* 72: 199–221.

Delobel, A. and C. D. Johnson (1998) First record of a seed-

- beetle on *Leucaena leucocephala* in West Africa. *Leucenet News* 5: 25–26.
- Delobel, A., G. Couturier, F. Kahn and J. A. Nilsson (1995a) Trophic relationships between palms and bruchids (Coleoptera, Bruchidae: Pachymerini) in Peruvian Amazonia. *Amazoniana* 13: 209–219.
- Delobel, A., H. Delobel, M. Tran, M. Sembène and S. H. Han (1995b) Observations sur les relations trophiques entre les bruches du genre *Carvedon* (Coléoptères, Bruchidae) et leurs plantes hôtes sauvages au Sénégal. *Bull. Inst. Fondam. Afriq. Noire Cheikh Anta Diop Dakar A* 48: 79–88.
- Delobel, B. and A. Delobel (2006) Dietary specialization in European species groups of seed beetles (Coleoptera: Bruchidae: Bruchinae). *Oecologia* 149: 428–443.
- Dongre, T. K., S. E. Pawar and M. R. Harwalkar (1993) Resistance to *Callosobruchus maculatus* (F.) (Coleoptera: Bruchidae) in pigeonpea (*Cajanus cajan* (L.) Millsp.) and other *Cajanus* species. *J. Stored Prod. Res.* 29: 319–322.
- Egorov, A. B. and M. E. Ter-Minassian (1983) *Bruchidae of East Siberia and Far East of the USSR*. Biol. Soil Stud. Inst. Akad. Nauk SSSR, Leningrad (in Russian).
- Ehrlich, P. R. and P. H. Raven (1964) Butterflies and plants: a study in coevolution. *Evolution* 18: 586–608.
- Elharith, E. A., U. T. Meulen and K. D. Gunther (1980) *Leucaena leucocephala*: a protein-rich and toxin containing fodder plant. *J. Anim. Physiol. Anim. Nut.* 44: 2–3.
- Farrell, B. D. (1998) “Inordinate fondness” explained: Why are there so many beetles? *Science* 281: 555–559.
- Farrell, B. D. and A. S. Sequeira (2004) Evolutionary rates in the adaptive radiation of beetles on plants. *Evolution* 58: 1984–2001.
- Fox, C. W. and T. A. Mousseau (1995) Determinants of clutch size and seed preference in a seed beetle, *Stator beali* (Coleoptera: Bruchidae). *Environ. Entomol.* 24: 1557–1561.
- Frenzel, M. and R. Brandl (1998) Diversity and composition of phytophagous insect guilds on Brassicaceae. *Oecologia* 113: 391–399.
- Fujii, K. and K. M. Wai (1990) Sex-ratio determination in three wasp species ectoparasitic on bean weevil larvae. In *Bruchids and Legumes: Economics, Ecology and Coevolution* (K. Fujii, A. M. R. Gatehouse, C. D. Johnson, R. Mitchel and T. Yoshida eds.). Kluwer, Dordrecht, pp. 331–340.
- Fujii, K., A. M. R. Gatehouse, C. D. Johnson, R. Mitchel and T. Yoshida (1990) *Bruchids and Legumes: Economics, Ecology and Coevolution*. Kluwer, Dordrecht, 407 pp.
- Futuyama, D. J. (1983) Selective factors in the evolution of host choice by phytophagous insects. In *Herbivorous Insects: Host Seeking Behavior and Mechanisms* (S. Ahmad ed.). Academic Press, New York, pp. 227–244.
- Henderson, L. (2001) *Alien Weeds and Invasive Plants. A Complete Guide to Declared Weeds and Invaders in South Africa*. Plant Protection Research Institute Handbook No. 12. Agricultural Research Council, Pretoria, 300 pp.
- Highley, E., E. J. Wright, H. J. Banks and B. R. Champ (1994) *Stored Product Protection: Proceeding of the 6th International Working Conference on Stored-Product Protection*. Vols. 1 and 2. CAB International, Wallingford, 1312 pp.
- Hossain, M. M., J. H. Brower and F. W. Tilton (1972) Sensitivity to an acute gamma radiation exposure of successively irradiated generations of cowpea weevil (Coleoptera: Bruchidae). *J. Econ. Entomol.* 65: 1566–1568.
- Huffaker, C. B. (1964) Fundamentals of biological weed control. In *Biological Control of Insect Pests and Weeds* (P. DeBach ed.). Chapman and Hall, London, pp. 631–649.
- Hughes, C. E. and C. D. Johnson (1996) New host records and notes on Bruchidae (Coleoptera) from *Leucaena Benth.* (Leguminosae, Mimosoideae) from Mexico, Central and South America. *J. Appl. Entomol.* 120: 137–141.
- Huis, A. van (1991) Biological methods of bruchid control in the tropics: a review. *Insect Sci. Appl.* 12: 87–102.
- Imura, O. (1990) Life histories of stored-product insects. In *Bruchids and Legumes: Economics, Ecology and Coevolution* (K. Fujii, A. M. R. Gatehouse, C. D. Johnson, R. Mitchel and T. Yoshida eds.). Kluwer, Dordrecht, pp. 257–269.
- Jaloux, B., A. Sanon, J. Huignard and J. P. Monge (2004) Interspecific relationships between the solitary ectoparasitoid, *Eupelmus vuilleti* (Craw.) (Eupelmidae), and its sympatric species, *Dinarmus basalis* (Rond.) (Pteromalidae), in the presence of their host, *Callosobruchus maculatus* Pic (Coleoptera Bruchidae). *J. Insect Behav.* 17: 793–808.
- Janzen, D. H. (1969) Seed-eaters versus seed size, number, toxicity and dispersal. *Evolution* 23: 1–27.
- Janzen, D. H. (1971) Seed predation by animals. *Annu. Rev. Ecol. Syst.* 2: 465–492.
- Janzen, D. H. (1977) How southern cowpea weevil larvae (Bruchidae: *Callosobruchus maculatus*) die on non-host seeds. *Ecology* 58: 921–927.
- Jerry, T. (1984) Evolution of insect host plant relationships. *Amer. Nat.* 124: 609–630.
- Jerry, T. and A. Szentesi (2003) Evolutionary aspects of host plant specialisation—a study on bruchids (Coleoptera: Bruchidae). *Oikos* 101: 207–215.
- Johnson, C. D. (1967) Notes on the systematics, host plants and bionomics of the Bruchid genera *Merobruchus* and *Stator*. *Pan-Pac. Entomol.* 43: 264–271.
- Johnson, C. D. (1970) Biosystematics of the Arizona, California and Oregon species of the seed beetle genus *Acanthoscelides* Schilsky (Coleoptera: Bruchidae). *Univ. Calif. Publ. Entomol.* 59: 1–116.
- Johnson, C. D. (1979) New host records for *Acanthoscelides* (Coleoptera: Bruchidae). *Pan-Pac. Entomol.* 55: 61–71.
- Johnson, C. D. (1981) Seed beetle host specificity and the systematics of the Leguminosae. In *Advances in Legume Systematics Part 2* (R. M. Polhill and P. H. Raven eds.). Royal Botanical Gardens, Kew, England, pp. 995–1027.
- Johnson, C. D. and D. H. Siemsen (1991) Expanded oviposition range by a seed beetle (Coleoptera: Bruchidae) in proximity to a normal host. *Environ. Entomol.* 20: 1577–1582.

- Johnson, C. D., B. J. Southgate and A. Delobel (2004) A revision of the Caryedontini (Coleoptera: Bruchidae: Pachymerinae) of Africa and the Middle East. *Mem. Amer. Entomol. Soc.* 44: 1–120.
- Julien, M. H. (1992) *Biological Control of Weeds: A World Catalogue of Agents and Their Target Weeds*. 3rd ed. CAB International, Wallingford. 196 pp.
- Kergoat, G. J., N. Alvarez, M. Hossaert-Mckey, N. Faure and J. F. Silvain (2005a) Parallels in the evolution of the two largest New and Old World seed-beetle genera (Coleoptera: Bruchidae). *Mol. Ecol.* 14: 4003–4021.
- Kergoat, G. J., A. Delobel, G. Fediere, B. Le Ru and J. F. Silvain (2005b) Both host-plant phylogeny and chemistry have shaped the African seed-beetle radiation. *Mol. Phylogenet. Evol.* 35: 602–611.
- Kergoat, G. J., J. F. Silvain, S. Buranapamichpan and M. Tuda (2007a) When insects help to resolve plant phylogeny: Evidence for a paraphyletic genus *Acacia* from the systematics and host-plant range of their seed-predators. *Zool. Scripta* 36: 143–152.
- Kergoat, G. J., J. F. Silvain, A. Delobel, M. Tuda and K.-W. Anton (2007b) Defining the limits of taxonomic conservatism in host-plant use for phytophagous insects: molecular systematics and evolution of host-plant associations in the seed-beetle genus *Bruchus* Linnaeus (Coleoptera: Chrysomelidae: Bruchinae). *Mol. Phylogenet. Evol.* 43: 251–269.
- Kingsolver, J. M. (1999) A new species of *Callosobruchus* (Coleoptera: Bruchidae) from Thailand and China. *Entomol. News* 110: 209–213.
- Kitch, I. W., R. E. Shade and L. L. Murdock (1991) Resistance to the cowpea weevil (*Callosobruchus maculatus*) larva in pods of cowpea (*Vigna unguiculata*). *Entomol. Exp. Appl.* 60: 183–192.
- Kiyoku, M. and R. Tsukuda (1968) Effect of <sup>137</sup>Cs on the sterilities of *Callosobruchus chinensis* (L.) and *Spodoptera litura* (F.). Studies on the ecology of insects sterilized artificially (gamma radiation). I. *Odokon-Chugoku* 10: 10–12 (in Japanese).
- Kobayashi, A., Y. Tanaka and M. Shimada (2003) Genetic variation of sex allocation in the parasitoid wasp *Heterospilus prosopidis*. *Evolution* 57: 2659–2664.
- Kondo, N., H. Yuasa and F. Maekawa (1987) *Resource-Handbook of Legumes*. Uchida Rokakuho Publ., Tokyo (in Japanese).
- Kondo, N., N. Ijichi, M. Shimada and T. Fukatsu (2002) Prevailing triple infection with *Wolbachia* in *Callosobruchus chinensis* (Coleoptera: Bruchidae). *Mol. Ecol.* 11: 167–180.
- Lambert, J. D. H. (1985) Bruchid control with traditionally used insecticidal plants *Hyptis spicigera* and *Cassia nigricans*. *Insect Sci. Appl.* 6: 167–170.
- Lewinsohn, T. M. (1991) Insects in flower heads of Asteraceae in southeast Brazil: a case study of tropical species richness. In *Plant-Animal Interactions: Evolutionary Ecology in Tropical and Temperate Regions* (P. W. Price, T. M. Lewinsohn, G. W. Fernandes and W. W. Benson eds.). Wiley, New York, pp. 525–559.
- Lingafelter, S. and J. Pakaluk (1997) Comments on the Bean beetle Chrysomelidae. *Chrysomela* 33: 3.
- Messina, F. J. (1984) Influence of cowpea pod maturity on the oviposition choices and larval survival of a bruchid beetle *Callosobruchus maculatus*. *Entomol. Exp. Appl.* 35: 241–248.
- Mitchell, R. (1977) Bruchid beetles and seed packaging by Palo Verde. *Ecology* 58: 644–651.
- Mitsunaga, T. and K. Fujii (1999) An experimental analysis of the relationship between species combination and community persistence. *Res. Popul. Ecol.* 41: 127–134.
- Moreno-Casasola, P., J. P. Grime and M. L. Martinez (1994) A comparative study of the effects of fluctuations in temperature and moisture supply on hard coat dormancy in seeds of coastal tropical legumes in Mexico. *J. Trop. Ecol.* 10: 67–86.
- Morimoto, K. (1990) A synopsis of the bruchid fauna of Japan. In *Bruchids and Legumes: Economics, Ecology and Coevolution* (K. Fujii, A. M. R. Gatchouse, C. D. Johnson, R. Mitchel and T. Yoshida eds.). Kluwer, Dordrecht, pp. 131–140.
- Morimoto, N. and K. Kiritani (1995) Fauna of exotic insects in Japan. *Bull. Natl. Inst. Agro-Environ. Sci.* 12: 87–120.
- Nakamura, H. (1966) The active-types observed in the adult of *Callosobruchus chinensis* L. *Jpn. J. Ecol.* 16: 236–241 (in Japanese with English summary).
- Neser, S. (1994) Conflicts of interest? The *Leucaena* controversy. *Plant Prot. News S. Afr.* 6: 8.
- Nilsson, J. A. and C. D. Johnson (1993) Laboratory hybridization of *Stator beali* and *S. limbatus*, with new host records for *S. limbatus* and *Mimosestes amicus*. *Southwest. Naturalist* 38: 385–387.
- Olckers, T. (2004) Targeting emerging weeds for biological control in South Africa: the benefit of halting the spread of alien plants at an early stage of their invasion. *S. Afr. J. Sci.* 100: 64–68.
- Oosthuizen, M. J. and U. W. Schmidh (1942) The toxicity of carbon dioxide to cowpea weevil. *J. Entomol. Soc. S. Afr.* 5: 55–110.
- Ouedraogo, A. P. and J. Huignard (1981) Polymorphism and ecological reactions in *Callosobruchus maculatus* F. (Coleoptera: Bruchidae) in Upper Volta. In *The Ecology of Bruchids Attacking Legumes (Pulses)* (V. Labeyrie ed.). Junk Publ., The Hague, pp. 175–184.
- Paynter, Q. E., S. V. Fowler, A. H. Goulay, M. L. Haines, H. M. Harman, S. R. Hona, P. G. Peterson, L. A. Smith, J. R. A. Wilson-Davey, C. J. Winks and T. M. Withers (2004) Safety in New Zealand weed biocontrol: a nationwide survey for impacts on non-target plants. *New Zeal. Pl. Prot.* 57: 102–107.
- Pfaffenberger, G. S. and C. D. Johnson (1976) Biosystematics of the first-stage larvae of some North American Bruchidae (Coleoptera). *U. S. Dep. Agr. Tech. Bull.* 1525: 1–75.
- Prevett, P. F. (1966) Observations on biology in the genus *Caryedon* Schoenherr (Coleoptera: Bruchidae) in Northern Nigeria with a list of associated parasitic Hymenoptera. *Proc. R. Entomol. Soc. Lond. A* 41: 9–16.
- Prevett, P. F. (1967) Notes on the biology, food plants and



- distribution of Nigerian Bruchidae with particular reference to the northern region. *Bull. Entomol. Soc. Nigeria* 1: 3-6.
- Templeton, M. F., J. L. Spencer and J. R. Miller (1991) Bean tumbling as a control measure for the common bean weevil, *Acanthoscelides obiectus*. *Entomol. Exp. Appl.* 60: 105-109.
- Teran, M. M. (1990) Some promising physical, botanical and chemical methods for the protection of grain legumes against bruchids in storage under Bangladesh conditions. In *Bruchids and Legumes. Economics, Ecology and Coevolution* (K. Fujii, A. M. R. Gatehouse, C. D. Johnson, R. Mitchel and T. Yoshida eds.). Kluwer, Dordrecht, pp. 63-73.
- Teddy, P. V. R., S. Todoriki, A. Miyano-shita, T. Imamura and T. Hayashi (2006) Effect of soft electron treatment on adzuki bean weevil, *Callosobruchus chinensis* (L.) (Col., Bruchidae). *J. Appl. Entomol.* 130: 393-399.
- Tingers, C. E. and J. C. Garrison (1975) Seed destruction in Indigobush *Amorpha* by a seed beetle. *J. Range Manag.* 28: 241-242.
- Tomero, J. and C. D. Johnson (2000) Revision of the genus *Zabrotex* Horn of Mexico (Coleoptera: Bruchidae: Amblycerinae). *Trans. Amer. Entomol. Soc.* 126: 221-274.
- Tsuno, I. (1967) Density effect and environmental temperature as the factors producing the active form of *Callosobruchus maculatus* (F.) (Coleoptera, Bruchidae). *J. Stored Prod. Res.* 2: 187-195.
- Satake, Y., H. Hara, S. Watari and T. Tominari (1989) *Wild Flowers of Japan: Woody Plants I*. Heibonsha, Tokyo.
- Schmale, I., F. L. Wickers, C. Cardona and S. Dorn (2001) Control potential of three hymenopteran parasitoid species against the bean weevil in stored beans: the effect of adult parasitoid nutrition on longevity and progeny production. *Biol. Control* 21: 134-139.
- Smith, C. W. (1985) Impact of alien plants on Hawaii's native biota. In *Hawaii's Terrestrial Ecosystems: Preservation and Management* (C. P. Stone and J. M. Scott eds.). Cooperative National Park Resources Studies Unit, University of Hawaii, Manoa, pp. 180-250.
- Southgate, B. J. (1979) Biology of the Bruchidae. *Annu. Rev. Entomol.* 24: 449-473.
- Steffan, J. R. (1981) The parasites of bruchids. In *The Ecology of Bruchids Attacking Legumes (Pulses)* (V. Labeyrie ed.). Junk Publ., The Hague, pp. 223-229.
- Sweetman, H. L. (1936) *The Biological Control of Insects: With a Chapter on Weed Control*. Comstock Publishing Co., Ithaca, NY. 461 pp.
- Szentesi, A. (1999) Predispersal seed predation of the introduced false indigo, *Amorpha fruticosa* L. in Hungary. *Acta Zool. Acad. Sci. Hung.* 45: 125-141.
- Szentesi, A. and T. Jermy (1995) Predispersal seed predation in leguminous species: seed morphology and bruchid distribution. *Oikos* 73: 23-32.
- Takano, M., Y. Toquenaga and K. Fujii (2001) Polymorphism of competition type and its genetics in *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Popul. Ecol.* 43: 265-273.
- Templeton, A. R. (1998) Nested clade analyses of phylogeographic data: testing hypotheses about gene flow and population history. *Mol. Ecol.* 7: 381-397.
- Templeton, A. R., K. A. Crandall and C. F. Sing (1992) A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping and DNA sequence data. 3. Cladogram estimation. *Genetics* 132: 619-633.
- Teran, A. L. (1962) Observaciones sobre Bruchidae del Noroeste Argentino. *Acta Zool. Lilloana* 18: 211-242.
- Toquenaga, Y. (1990) The mechanisms of contest and scramble competition in bruchid species. In *Bruchids and Legumes: Economics, Ecology and Coevolution* (K. Fujii, A. M. R. Gatehouse, C. D. Johnson, R. Mitchel and T. Yoshida eds.). Kluwer, Dordrecht, pp. 341-349.
- Tuda, M. (1997) Coexistence and evolution in experimental ecological systems. In *Mathematical Ecology* (Y. Iwasa ed.). Kyoritsu Shuppan, Tokyo, pp. 69-79 (in Japanese).
- Tuda, M. (1998) Evolutionary character changes and population responses in an insect host-parasitoid experimental system. *Res. Popul. Ecol.* 40: 293-299.
- Tuda, M. (2003) A new species of *Callosobruchus* (Coleoptera: Bruchidae) feeding on seeds of *Dunbaria* (Fabaceae), a closely related species to a stored-bean pest, *C. chinensis*. *Appl. Entomol. Zool.* 38: 197-201.
- Tuda, M. and Y. Iwasa (1998) Evolution of contest competition and its effect on host-parasitoid dynamics. *Evol. Ecol.* 12: 855-870.
- Tuda, M. and K. Morimoto (2004) A new species *Megabruchidius sophorae* (Coleoptera, Bruchidae), feeding on seeds of *Styphnolobium* (Fabaceae) new to Bruchidae. *Zool. Sci.* 21: 105-110.
- Tuda, M. and M. Shimada (2005) Complexity, evolution, and persistence in host-parasitoid experimental systems with *Callosobruchus* beetles as the host. *Adv. Ecol. Res.* 37: 37-75.
- Tuda, M., T. Fukatsu and M. Shimada (1995) Species differentiation of bruchid beetles (Coleoptera: Bruchidae) analyzed by mitochondrial DNA polymorphism. *Appl. Entomol. Zool.* 30: 377-380.
- Tuda, M., K. Shima, C. D. Johnson and K. Morimoto (2001) Establishment of *Acanthoscelides pallidipennis* (Coleoptera: Bruchidae) feeding in seeds of the introduced legume *Amorpha fruticosa*, with a new record of its *Eupelmus* parasitoid in Japan. *Appl. Entomol. Zool.* 36: 269-276.
- Tuda, M., N. Wasano, N. Kondo, S.-B. Horng, L.-Y. Chou and Y. Tateishi (2004) Habitat-related mtDNA polymorphism in the stored-bean pest *Callosobruchus chinensis* (Coleoptera: Bruchidae). *Bull. Entomol. Res.* 94: 75-80.
- Tuda, M., L.-Y. Chou, C. Niyomdham, S. Buranapanichpan and Y. Tateishi (2005) Ecological factors associated with pest status in *Callosobruchus* (Coleoptera: Bruchidae): high host specificity of non-pests to Cajaninae (Fabaceae). *J. Stored Prod. Res.* 41: 31-45.
- Tuda, M., J. Ronn, S. Buranapanichpan, N. Wasano and G. Arnqvist (2006) Evolutionary diversification of the bean beetle genus *Callosobruchus* (Coleoptera: Bruchidae): traits associated with stored-product pest status.

*Mol. Ecol.* 15: 3541–3551.

Udayagiri, S. and S. R. Wadhi (1989) *Catalog of Bruchidae*. American Entomological Institute, Gainesville. 301 pp.

UNEP (2000) *The Montreal Protocol on Substances That Deplete the Ozone Layer*. UNOP, Kenya.

Utida, S. (1954) "Phase" dimorphism observed in the laboratory population of the cowpea weevil, *Callosobruchus quadrimaculatus*. *J. Appl. Zool.* 18: 161–168 (in Japanese with English summary).

Utida, S. (1981) Polymorphism and phase dimorphism in *Callosobruchus*. In *The Ecology of Bruchids Attacking Legumes (Pulses)* (V. Labeyrie ed.). Dr W. Junk Publishers, The Hague, pp. 143–147.

Watanabe, N. (1985) Oviposition habit of *Sulcobruchus sauteri* (Pic) and its significance in speculation on the pre-agricultural life of seed beetles attacking stored pulses (Coleoptera, Bruchidae). *Kontyu* 53: 391–397.

Wu, S.-H., S.-M. Chaw and M. Rejmánek (2003) Naturalized Fabaceae (Leguminosae) species in Taiwan: the first

approximation. *Bot. Bull. Acad. Sin.* 44: 59–66.

Wu, Z., K. R. Hopper, P. J. Ode, R. W. Fuester, M. Tuda and G. E. Heimpel (2005) Single-locus complementary sex determination absent in *Heterospilus prosopidis* (Hymenoptera: Braconidae). *Heredity* 95: 228–234.

Yoneda, M., M. Kaneda and H. Akiyama (1990) Plant quarantine and fumigation of imported grain legumes in Japan. In *Bruchids and Legumes: Economics, Ecology and Coevolution* (K. Fujii, A. M. R. Gatehouse, C. D. Johnson, R. Mitchel and T. Yoshida eds.). Kluwer, Dordrecht, pp. 115–120.

Yoshida, T. (1990) Historical review of bruchid studies in Japan. In *Bruchids and Legumes: Economics, Ecology and Coevolution* (K. Fujii, A. M. R. Gatehouse, C. D. Johnson, R. Mitchel and T. Yoshida eds.). Kluwer, Dordrecht, pp. 1–24.

Zacher, F. (1952) Die Nahrungspflanzen der samenkafer. *Zeits. Angew. Entomol.* 33: 460–480.